

Bibliografia

Alavi FK, Zawada ET, Simmons JL, 1995: Renal hemodynamic and histological consequences of diets high in unsaturated fat, protein or sucrose in obese Zucker rats. *Clin Nephrol* 43:122-130.

Alexy U, Remer T, Manz F, Neu CM, Schoenau E, 2005: Long-term protein intake and dietary potential renal acid load are associated with bone remodeling and remodeling at the proximal radius in healthy children. *Am J Clin Nutr* 82:1107–1114

Allen LH, Oddoye EA, Margen S, 1979: Protein-induced hypercalciuria: A longer term study. *Am J Clin Nutr* 32:741- 749.

Arnadottir M, Hultberg B, Nilsson-Ehle P, Thysell H, 1996: The effect of reduced glomerular filtration rate on plasma total homocysteine concentration. *Scand J Clin Lab Invest* 56:41-46.

Amanzadeh J, Gitomer WL, Zerwekh JE, et al, 2003: Effect of high protein diet on stone-forming propensity and bone loss in rats. *Kidney Int* 64:2142-2149.

Amore A, Coppo R, Roccatello D, et al, 1985: Single kidney function: Effect of acute protein and water loading on microalbuminuria. *Am J Med* 84:711-716.

Appel LJ, Moore TJ, Obarzanek E, et al, 1997: A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *N Engl J Med* 336: 1117-1124.

Apfelbaum M, Baigts F, Giachetti I, Serog P, 1981: Effects of a high protein very-low-energy diet on ambulatory subjects with special reference to nitrogen balance. *Int J Obes* 5:117-130.

Arroyave G, Wilson D, Behar M, Scrimshaw N, 1961: Serum and urinary creatinine in children with severe protein malnutrition. *Am J Clin Nutr* 9:176-179.

Atkins R, 1999: Dr Atkins' New Diet Revolution. New York, NY, Avon.

Azar G, Bloom W, 1963: Similarities of carbohydrate deficiency and fasting. II. Ketones, nonesterified fatty acids, and nitrogen excretion. *Arch Intern Med* 112:338-343.

Ball D, Maughan RJ, 1997: Blood and urine acid-base status of premenopausal omnivorous and vegetarian women. *Br J Nutr* 78:683-693.

Benoit F, Martin R, Watten R, 1965: Changes in body composition during weight reduction in obesity. *Ann Intern Med* 63:604-612.

Bergstrom J, Ahlberg M, Alvestrand A, 1985: Influence of protein intake on renal hemodynamics and plasma hormone concentrations in normal subjects. *Acta Med Scand* 217:189- 196.

Bilo HJ, Schaap GH, Blaak E, Gans RO, Oe PL, Donker AJ, 1989: Effects of chronic and acute protein administration on renal function in patients with chronic renal insufficiency. *Nephron* 53:181-187.

Bloom W, 1962: Inhibition of salt excretion by carbohydrate. *Arch Intern Med* 109:25-32.

Bloom W, Azar G, 1963: Similarities of carbohydrate deficiency and fasting. I. Weight loss, electrolyte excretion, and fatigue. *Arch Intern Med* 112:333-337.

Bohlender JM, Franke S, Stein G, Wolf G, 2005: Advanced glycation end products and the kidney. *Am J Physiol Renal Physiol.* 289(4), F645-59.

Borghi L, Schianchi T, Meschi T, et al, 2002: Comparison of two diets for the prevention of recurrent stones in idiopathic hypercalciuria. *N Engl J Med* 346:77-84.

Bosch JP, Saccaggi A, Lauer A, Ronco C, Belledonne M, Glabman S, 1983: Renal functional reserve in humans. Effect of protein intake on glomerular filtration rate. *Am J Med* 75:943-950.

Bowen J, Noakes M, Clifton PM, 2004: A high dairy protein, high-calcium diet minimizes bone turnover in overweight adults during weight loss. *J Nutr* 134:568-573.

Brandle E, Hesse A, 1993: Influence of oral protein intake on the calcium excretion and the glomerular filtration rate. *Urol Res* 21:153A.

Brandle E, Sieberth HG, Hautmann RE, 1996: Effect of chronic dietary protein intake on the renal function in healthy subjects. *Eur J Clin Nutr* 50:734-740.

Bravata DM, Sanders L, Huang J, Krumholz HM, Olkin I, Gardner CD, 2003: Efficacy and safety of low-carbohydrate diets: A systematic review. *JAMA* 289:1837-1850.

Brehm BJ, Seeley RJ, Daniels SR, D'Alessio DA, 2003: A randomized trial comparing a very low carbohydrate diet and a calorie-restricted low fat diet on body weight and cardiovascular risk factors in healthy women. *J Clin Endocrinol Metab* 88:1617-1623.

Brenner BM, Meyer TW, Hostetter TH, 1982: Dietary protein intake and the progressive nature of kidney disease: The role of hemodynamically

mediated glomerular injury in the pathogenesis of progressive glomerular sclerosis in aging, renal ablation, and intrinsic renal disease. *N Engl J Med* 307:652-659.

Breslau NA, Brinkley L, Hill KD, Pak CY, 1988: Relationship of animal protein-rich diet to kidney stone formation and calcium metabolism. *J Clin Endocrinol Metab* 66:140- 146.

Brochner-Mortensen J, Rickers H, Balslev I, 1980: Renal function and body composition before and after intestinal bypass operation in obese patients. *Scand J Clin Lab Invest* 40:695-702.

Carmo I, Santos O, Camolas J, Vieira J, Carreira M, Medina L et al, 2006: Prevalence of obesity in Portugal. *Obes Rev.* 7(3):233-7.

Carmo I, Santos O, Camolas J, Vieira J, Carreira M, Medina L et al, 2008: Overweight and obesity in Portugal: national prevalence in 2003-2005. *Obes Rev.* 9(1): 11-9. Epub 2007 Nov 23.

Castellino P, DeFronzo RA, 1988: Effect of plasma amino acid and hormone concentrations on renal plasma flow and glomerular filtration rate. *Blood Purif* 6:240-249.

Chagnac A, Weinstein T, Korzets A, Ramadan E, Hirsch J, Gafter U,, 2000: Glomerular hemodynamics in severe obesity. *Am J Physiol Renal Physiol* 278:F817-F822.

Chagnac A, Weinstein T, Herman M, Hirsh J, Gafter U, Ori Y 2003: The effects of weight loss on renal function in patients with severe obesity. *J Am Soc Nephrol* 14:1480-1486.

ChanAY, Cheng ML, Keil LC, Myers BD, 1988: Functional response of healthy and diseased glomeruli to a large, protein-rich meal. *J Clin Invest* 81:245-254.

Choi HK, Atkinson K, Karlson EW, Willett W, Curhan G, 2004: Purine-rich foods, dairy and protein intake, and the risk of gout in men. *N Engl J Med* 350:1093-1103.

Claris-Appiani A, Assael BM, Tirelli AS, Marra G, Cavanna G, 1988: Lack of glomerular hemodynamic stimulation after infusion of branched-chain amino acids. *Kidney Int* 33:91-94.

Coppo R, Amore A, Roccatello D, et al, 1988: Microalbuminuria in single kidney patients: Relationship with protein intake. *Clin Nephrol* 29:219-228.

Curhan GC, Willett WC, Rimm EB, Stampfer MJ, 1993: A prospective study of dietary calcium and other nutrients and the risk of symptomatic kidney stones. *N Engl J Med* 328:833-838.

Curhan GC, Willett WC, Knight EL, Stampfer MJ, 2004: Dietary factors and the risk of incident kidney stones in younger women: Nurses' Health Study II. *Arch Intern Med* 164:885-891.

DeHaven J, Sherwin R, Hendler R, Felig P 1980: Nitrogen and sodium balance and sympathetic-nervous-system activity in obese subjects treated with a low-calorie protein or mixed diet. *N Engl J Med* 302:477-482.

Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids. Washington, DC, Institute of Medicine, 2002, p S16

Eisenstein J, Roberts SB, Dallal G, Saltzman E, 2002: High-protein weight-loss diets: Are they safe and do they work? A review of the experimental and epidemiologic data. *Nutr Rev* 60:189-200.

Feskanich D, Willett WC, Stampfer MJ, Colditz GA, 1996: Protein consumption and bone fractures in women. *Am J Epidemiol* 143:472-479.

Fried LF, Orchard TJ, Kasiske BL, 2001: Effect of lipid reduction on the progression of renal disease: A metaanalysis. *Kidney Int* 59:260-269.

Friedman AN, Bostom AG, Selhub J, Levey AS, Rosenberg IH, 2001: The kidney and homocysteine metabolism. *JAm Soc Nephrol* 12:2181-2189.

Foster GD, Wyatt HR, Hill JO, et al, 2003: A randomized trial of a low-carbohydrate diet for obesity. *N Engl J Med* 348:2082-2090.

Gambaro G, Favaro S, D'Angelo A, 2001: Risk for renal failure in nephrolithiasis. *Am J Kidney Dis* 37:233-243.

Gerstein HC, Mann JF, Yi Q, et al, 2001: Albuminuria and risk of cardiovascular events, death, and heart failure in diabetic and nondiabetic individuals. *JAMA* 286:421-426.

Giordano M, Castellino P, McConnell EL, DeFronzo RA, 1994: Effect of amino acid infusion on renal hemodynamics in humans: A dose-response study. *Am J Physiol* 267:F703- F708.

Giovannetti S, Maggiore Q, 1964: A low-nitrogen diet with proteins of high biological value for severe chronic uraemia. *Lancet* 37:1000-1003.

Ginty F, 2003: Dietary protein and bone health. *Proc Nutr Soc* 62:867-876.

Graf H, Stummvoll HK, Luger A, Prager R, 1983: Effect of amino acid infusion on glomerular filtration rate. *N Engl J Med* 308:159-160.

Greenhaff PL, Gleeson M, Maughan RJ, 1988: Diet-induced metabolic acidosis and the performance of high intensity exercise in man. *Eur J Appl Physiol Occup Physiol* 57:583-590.

Goldberg A, Guggenheim K, 1962: The digestive release of amino acids and their concentrations in the portal plasma of rats after protein feeding. *Biochem J* 83:129-135.

Hammond KA, Janes DN, 1998: The effects of increased protein intake on kidney size and function. *J Exp Biol* 201:2081-2090.

Haulrik N, Toubro S, Dyerberg J, Stender S, Skov AR, Astrup A, 2002: Effect of protein and methionine intakes on plasma homocysteine concentrations: A 6-mo randomized controlled trial in overweight subjects. *Am J Clin Nutr* 76:1202-1206.

Hegsted M, Linkswiler HM, 1981: Long-term effects of level of protein intake on calcium metabolism in young adult women. *J Nutr* 111:244-251.

Hegsted M, Schuette SA, Zemel MB, Linkswiler HM, 1981: Urinary calcium and calcium balance in young men as affected by level of protein and phosphorus intake. *J Nutr* 111:553-562.

Henegar JR, Bigler SA, Henegar LK, Tyagi SC, Hall JE, 2001: Functional and structural changes in the kidney in the early stages of obesity. *J Am Soc Nephrol* 12:1211-1217.

Hiatt RA, Ettinger B, Caan B, Quesenberry CP Jr, Duncan D, Citron JT, 1996: Randomized controlled trial of a low animal protein, high fiber diet in the prevention of recurrent calcium oxalate kidney stones. *Am J Epidemiol* 144:25-33.

Hillege HL, Fidler V, Diercks GF, et al, 2002: Urinary albumin excretion predicts cardiovascular and noncardiovascular mortality in general population. *Circulation* 106:1777- 1782.

Hoogeveen EK, Kostense PJ, Jager A, et al, 1998: Serum homocysteine level and protein intake are related to risk of microalbuminuria: The Hoorn Study. *Kidney Int* 54:203- 209.

Ichikawa I, Purkerson ML, Klahr S, Troy JL, Martinez- Maldonado M, Brenner BM: Mechanism of reduced glomerular filtration rate in chronic malnutrition. *J Clin Invest* 65:982-988, 1980

Ihle BU, Becker GJ, Whitworth JA, Charlwood RA, Kincaid-Smith PS, 1989: The effect of protein restriction on the progression of renal insufficiency. *N Engl J Med* 321:1773- 1777.

Jacques PF, Bostom AG, Wilson PW, Rich S, Rosenberg IH, Selhub J, 2001: Determinants of plasma total homocysteine concentration in the Framingham offspring cohort. *Am J Clin Nutr* 73:613-621.

Johnson RJ, Kivlighn SD, Kim YG, Suga S, Fogo AB, 1999: Reappraisal of the pathogenesis and consequences of hyperuricemia in hypertension, cardiovascular disease, and renal disease. *Am J Kidney Dis* 33:225-234.

Johnston CS, Tjonn SL, Swan PD, White A, Sears B, 2006: Low-carbohydrate, high-protein diets that restrict potassium-rich fruits and vegetables promote calciuria *Osteoporos Int* 17:1820–1821.

Kalk WJ, Osler C, Constable J, Kruger M, Panz V, 1992: Influence of dietary protein on glomerular filtration and urinary albumin excretion in insulin-dependent diabetes. *Am J Clin Nutr* 56:169-173.

Kark R, Johnson R, Lewis J, 1945: Defects of pemmican as an emergency ration for infantry troops. *War Med* 7:345- 352.

Kasike BL, Cleary MP, O'Donnell MP, Keane WF, 1986: Effects of carbohydrate restriction on renal injury in the obese Zucker rat. *Am J Clin Nutr* 44:56-65.

Kasike BL, Lakatua JD, Ma JZ, Louis TA, 1998: A meta-analysis of the effects of dietary protein restriction on the rate of decline in renal function. *Am J Kidney Dis* 31:954- 961.

Kaysen GA, Gambertoglio J, Jimenez I, Jones H, Hutchison FN, 1986: Effect of dietary protein intake on albumin homeostasis in nephrotic patients. *Kidney Int* 29:572-577.

Kerstetter JE, Mitnick ME, Gundberg CM, et al, 1999: Changes in bone turnover in young women consuming different levels of dietary protein. *J Clin Endocrinol Metab* 84:1052-1055.

Klahr S, Levey AS, Beck GJ, et al, 1994: The effects of dietary protein restriction and blood-pressure control on the progression of chronic renal disease. Modification of Diet in Renal Disease Study Group. *N Engl J Med* 330:877-884.

Kok DJ, Iestra JA, Doorenbos CJ, Papapoulos SE, 1990: The effects of dietary excesses in animal protein and in sodium on the composition and the crystallization kinetics of calcium oxalate monohydrate in urines of healthy men. *J Clin Endocrinol Metab* 71:861-867.

Kontessis P, Jones S, Dodds R, et al 1990: Renal, metabolic and hormonal responses to ingestion of animal and vegetable proteins. *Kidney Int* 38:136-144.

Knight EL, Stampfer MJ, Hankinson SE, Spiegelman D, Curhan GC, 2003: The impact of protein intake on renal function decline in women with normal renal function or mild renal insufficiency. *Ann Intern Med* 138:460-467.

KeaneWF, Kasiske BL, O'Donnell MP, 1988: Hyperlipidemia and the progression of renal disease. *Am J Clin Nutr* 47:157-160.

Lacroix M, Gaudichon C, Martin A, et al 2004: A longterm high-protein diet markedly reduces adipose tissue without major side-effects in Wistar male rats. *Am J Physiol Regul Integr Comp Physiol* 287:934-942.

Lee KE, Summerill RA, 1982: Glomerular filtration rate following administration of individual amino acids in conscious dogs. *Q J Exp Physiol* 67:459-465.

Lemann J Jr, Gray RW, Maierhofer WJ, Cheung HS, 1986: The importance of renal net acid excretion as a determinant of fasting urinary calcium excretion. *Kidney Int* 29:743-746.

Levinsky N, Berliner R, 1959: The role of urea in the urine concentrating mechanism. *J Clin Invest* 38:741-748.

Lin PH, Ginty F, Appel LJ, et al, 2003: The DASH diet and sodium reduction improve markers of bone turnover and calcium metabolism in adults. *J Nutr* 133:3130-3136.

Locatelli F, Alberti D, Graziani G, Bucciatti G, Redaelli B, Giangrande A, 1991: Prospective, randomised, multicentre trial of effect of protein restriction on progression of chronic renal insufficiency. Northern Italian Cooperative Study Group. *Lancet* 337:1299-1304.

Luft FC, Fineberg NS, Sloan RS, Hunt JN, 1983: The effect of dietary sodium and protein on urine volume and water intake. *J Lab Clin Med* 101:605-610.

de Luis DA, Fernandez N, Aller R, De Luis J, Arranz M, Izaola O, 2003: Relation between total homocysteine levels and beer intake in patients with diabetes mellitus type 2. *Ann Nutr Metab* 47:119-123.

Luscombe ND, Clifton PM, Noakes M, Farnsworth E, Wittert G 2003: Effect of a high-protein, energy-restricted diet on weight loss and energy expenditure after weight stabilization in hyperinsulinemic subjects. *Int J Obes Relat Metab Disord* 27:582-590.

MacKay E, MacKay L, Addis T, 1928: Factors which determine renal weight. V. The protein intake. *Am J Physiol* 86:459-470.

Metcalf PA, Baker JR, Scragg RK, Dryson E, Scott AJ, Wild CJ, 1993: Dietary nutrient intakes and slight albuminuria in people at least 40 years old. *Clin Chem* 39:2191-2198.

Mimran A, Ribstein J, DuCailar G, Halimi JM, 1994: Albuminuria in normals and essential hypertension. *J Diabetes Complications* 8:150-156.

Mitch WE, Walser M, Steinman TI, Hill S, Zeger S, Tungsanga K, 1984: The effect of a keto acid-amino acid supplement to a restricted diet on the progression of chronic renal failure. *N Engl J Med* 311:623-629.

Munger RG, Cerhan JR, Chiu BC, 1999: Prospective study of dietary protein intake and risk of hip fracture in postmenopausal women. *Am J Clin Nutr* 69:147-152.

National Kidney Foundation, 2002: K/DOQI Clinical Practice Guidelines for Chronic Kidney Disease: Evaluation, Classification, and Stratification. *Am J Kidney Dis* 39:S1- S266, (suppl 1)

New SA, Robins SP, Campbell MK, et al, 2000: Dietary influences on bone mass and bone metabolism: Further evidence of a positive link between fruit and vegetable consumption and bone health? *Am J Clin Nutr* 71:142-151.

O'Connor WJ, Summerill RA, 1976: The effect of a meal of meat on glomerular filtration rate in dogs at normal urine flows. *J Physiol* 256:81-91.

Pedrini MT, Levey AS, Lau J, Chalmers TC, Wang PH, 1996: The effect of dietary protein restriction on the progression of diabetic and nondiabetic renal diseases: A metaanalysis. *Ann Intern Med* 124:627-632.

Pilkington T, Gainsborough H, Rosenoer C, Carey M, 1960: Diet and weight-reduction in the obese. *Lancet* 16:856-858.

Pitts R, 1944: The effects of infusing glycine and of varying the dietary protein intake on renal hemodynamics in the dog. *Am J Physiol* 142:355-365.

Premen AJ, 1989: Nature of the renal hemodynamic action of amino acids in dogs. *Am J Physiol* 256:F516-F523.

Pullman T, Alving A, Dern R, Landowne M, 1954: The influence of dietary protein intake on specific renal functions in normal man. *J Lab Clin Med* 44:320-332

Reddy ST, Wang CY, Sakhaee K, Brinkley L, Pak CY, 2002: Effect of low-carbohydrate high-protein diets on acidbase balance, stone-forming propensity, and calcium metabolism. *Am J Kidney Dis* 40:265-274.

Reisin E, Abel R, Modan M, Silverberg DS, Eliahou HE, Modan B, 1978: Effect of weight loss without salt restriction on the reduction of blood pressure in overweight hypertensive patients. *N Engl J Med* 298:1-6.

Ribstein J, du Cailar G, Mimran A, 1995: Combined renal effects of overweight and hypertension. *Hypertension* 26:610- 615.

Riley MD, Dwyer T, 1998: Microalbuminuria is positively associated with usual dietary saturated fat intake and negatively associated with usual dietary protein intake in people with insulin-dependent diabetes mellitus. *Am J Clin Nutr* 67:50-57.

Rodriguez-Iturbe B, Herrera J, Garcia R, 1988: Relationship between glomerular filtration rate and renal blood flow at different levels of protein-induced hyperfiltration in man. *Clin Sci (Colch)* 74:11-15.

Russo LM, Bakris GL, Comper WD, 2002: Renal handling of albumin: A critical review of basic concepts and perspective. *Am J Kidney Dis* 39:899-919.

Sacks FM, Svetkey LP, Vollmer WM, et al, 2001: Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. DASH Sodium Collaborative Research Group. *N Engl J Med* 344:3- 10.

Samaha FF, Iqbal N, Seshadri P, et al, 2003: A lowcarbohydrate as compared with a low-fat diet in severe obesity. *N Engl J Med* 348:2074-2081.

Scaglione R, Ganguzza A, Corrao S, et al, 1995: Central obesity and hypertension: Pathophysiologic role of renal haemodynamics and function. *Int J Obes Relat Metab Disord* 19:403-409.

Schaap GH, Bilo HJ, Alferink TH, Oe PL, Donker AJ, 1987: The effect of a high protein intake on renal function of patients with chronic renal insufficiency. *Nephron* 47:1-6.

Selhub J, Miller JW, 1992: The pathogenesis of homocysteinemia: Interruption of the coordinate regulation by Sadenosylmethionine of the remethylation and transsulfuration of homocysteine. *Am J Clin Nutr* 55:131-138.

Seney FD Jr, Persson EG, Wright FS 1987: Modification of tubuloglomerular feedback signal by dietary protein. *Am J Physiol* 252:F83-F90.

Sigler M, 1975: The mechanism of the natriuresis in fasting. *J Clin Invest* 55:377-387.

Skov AR, Toubro S, Ronn B, Holm L, Astrup A, 1999: Randomized trial on protein vs carbohydrate in ad libitum fat reduced diet for the treatment of obesity. *Int J Obes Relat Metab Disord* 23:528-536.

Skov AR, Toubro S, Bulow J, Krabbe K, Parving HH, Astrup A, 1999: Changes in renal function during weight loss induced by high vs low-protein low-fat diets in overweight subjects. *Int J Obes Relat Metab Disord* 23:1170-1177.

Smit E, Nieto FJ, Crespo CJ, Mitchell P, 1999: Estimates of animal and plant protein intake in US adults: Results from the Third National Health and Nutrition Examination Survey, 1988-1991. *J Am Diet Assoc* 99:813-820.

Solerte SB, Rondanelli M, Giaccherio R, et al, 1999: Serum glucagon concentration and hyperinsulinaemia influence renal haemodynamics and urinary protein loss in normotensive patients with central obesity. *Int J Obes Relat Metab Disord* 23:997-1003.

Solling K, Christensen CK, Solling J, Christiansen JS, Mogensen CE, 1986: Effect on renal haemodynamics, glomerular filtration rate and albumin excretion of high oral protein load. *Scand J Clin Lab Invest* 46:351-357.

St Jeor ST, Howard BV, Prewitt TE, Bovee V, Bazzarre T, Eckel RH, 2001: Dietary protein and weight reduction: A statement for healthcare professionals from the Nutrition Committee of the Council on Nutrition, Physical Activity, and Metabolism of the American Heart Association. *Circulation* 104:1869-1874.

Stern L, Iqbal N, Seshadri P, et al, 2004: The effects of low-carbohydrate versus conventional weight loss diets in severely obese adults: One-year follow-up of a randomized trial. *Ann Intern Med* 140:778-785.

Stolzenberg-Solomon RZ, Miller ER III, Maguire MG, Selhub J, Appel LJ, 1999: Association of dietary protein intake and coffee consumption with serum homocysteine concentrations in an older population. *Am J Clin Nutr* 69:467-475.

Sutton RA, Wong NL, Dirks JH, 1979: Effects of metabolic acidosis and alkalosis on sodium and calcium transport in the dog kidney. *Kidney Int* 15:520-533.

Talbott JH, Terplan KL, 1960: The kidney in gout. *Medicine* (Baltimore) 39:405-467.

Toeller M, Buyken A, Heitkamp G, et al, 1997: Protein intake and urinary albumin excretion rates in the EURODIAB IDDM Complications Study. *Diabetologia* 40:1219- 1226.

Uribarri J, Tuttle KR, 2006: Advanced Glycation End Products and Nephrotoxicity of High-Protein Diets. *Clin J Am Soc Nephrol* 1: 1293–1299.

US Renal Data System, 2004: USRDS 2003 Annual Data Report. The National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD.

Valensi P, Assayag M, Busby M, Paries J, Lormeau B, Attali JR, 1996: Microalbuminuria in obese patients with or without hypertension. *Int J Obes Relat Metab Disord* 20:574- 579.

Verhoef P, van Vliet T, Olthof MR, Katan MB, 2005: A high-protein diet increases postprandial but not fasting plasma total homocysteine concentrations: a dietary controlled, crossover trial in healthy volunteers., *Am J Clin Nutr.* 82(3):553-8.

Viberti G, Boggetti E, Wiseman MJ, Dodds R, Gross JL, Keen H, 1987: Effect of protein-restricted diet on renal response to a meat meal in humans. *Am J Physiol* 253:F388- F393.

Wagner EA, Falciglia GA, Amlal H, Levin L, Soleimani M, 2007: Short-Term Exposure to a High-Protein Diet Differentially Affects Glomerular Filtration Rate but Not Acid-Base Balance in Older Compared to Younger Adults. *J Am Diet Assoc.* 107:1404-1408.

Ward M, McNulty H, Pentieva K, et al, 2000: Fluctuations in dietary methionine intake do not alter plasma homocysteine concentration in healthy men. *J Nutr* 130:2653-2657.

Westman EC, Yancy WS, Edman JS, Tomlin KF, Perkins CE, 2002: Effect of 6-month adherence to a very low carbohydrate diet program. *Am J Med* 113:30-36.

Wetzels JF, Hoitsma AJ, Berden JH, Koene , 1988: Renal hemodynamic effects of a short-term high protein and low protein diet in patients with renal disease. *Clin Nephrol* 30:42-47.

Wiseman MJ, Hunt R, Goodwin A, Gross JL, Keen H, Viberti GC, 1987: Dietary composition and renal function in healthy subjects. *Nephron* 46:37-42.

Woods LL, 1993: Mechanisms of renal hemodynamic regulation in response to protein feeding. *Kidney Int* 44:659-675.

Wright JD, Kennedy-Stephanson J, Wang CY, Mc- Dowell MA, Johnson CI, 2004: Trends in intake of energy and macronutrients—United States, 1971-2000. *MMWR Morb Mortal Wkly Rep* 53:80-82.

Wrone EM, Carnethon MR, Palaniappan L, Fortmann SP, 2003: Association of dietary protein intake and microalbuminuria in healthy adults: Third National Health and Nutrition Examination Survey. *Am J Kidney Dis* 41:580-587.

Yancy WS Jr, Olsen MK, Guyton JR, Bakst RP, Westman EC, 2004: A low-carbohydrate, ketogenic diet versus a low-fat diet to treat obesity and hyperlipidemia: A randomized, controlled trial. *Ann Intern Med* 140:769-777.

Yang M, Itallie TV, 1976: Composition of weight lost during short-term weight reduction. *J Clin Invest* 58:722- 730.

Yuyun MF, Khaw KT, Luben R, et al, 2004: A prospective study of microalbuminuria and incident coronary heart disease and its prognostic significance in a British population: The EPIC-Norfolk Study. *Am J Epidemiol* 159:284-293.

Zemel MB, Schuette SA, Hegsted M, Linkswiler HM, 1981: Role of the sulfur-containing amino acids in proteininduced hypercalciuria in men. *J Nutr* 111:545-552.